

Pediatric Hypotension / Shock

History

- Blood loss
- Fluid loss
- Vomiting
- Diarrhea
- Fever
- Infection

Signs and Symptoms


- Restlessness, confusion, weakness
- Dizziness
- Tachycardia
- Hypotension (Late sign)
- Pale, cool, clammy skin
- Delayed capillary refill
- Dark-tarry stools

Differential

- Shock
 - Hypovolemic
 - Cardiogenic
 - Septic
 - Neurogenic
 - Anaphylactic
- Trauma
- Infection
- Dehydration
- Congenital heart disease
- Medication or Toxin

	Blood Glucose Analysis Procedure
A	IV / IO Procedure
P	Cardiac Monitor
	Pediatric Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol PM 2 <i>if indicated</i>

History and Exam Suggest Type of Shock



Hypotension / Shock

Norepinephrine 0.05 - 0.1 mcg/kg/min titrate up to 2 mcg/kg

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Lowest blood pressure by age: < 31 days: > 60 mmHg. 31 days to 1 year: > 70 mmHg. Greater than 1 year: 70 + 2 x age in years.**
- **Consider all possible causes of shock and treat per appropriate protocol. Majority of decompensation in pediatrics is airway related.**
- **Decreasing heart rate and hypotension occur late in children and are signs of imminent cardiac arrest.**
- **Shock may be present with a normal blood pressure initially.**
- **Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.**
- **Consider all possible causes of shock and treat per appropriate protocol.**
- **Hypovolemic Shock:**
 - Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding.
- **Cardiogenic Shock:**
 - Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricular / septum / valve / toxins.
- **Distributive Shock:**
 - Septic
 - Anaphylactic
 - Neurogenic: Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.
 - Toxic
- **Obstructive Shock:**
 - Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.
 - Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.
- **Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:**
 - Body cannot produce enough steroids (glucocorticoids / mineralocorticoids.) May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate. Usually hypotensive with nausea, vomiting, dehydration and / or abdominal pain. **If suspected Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list. May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.**